

MAT8201, Algebraic Topology

Assignment 2

Due in-class on Tuesday, March 5

Numbered exercises are from Hatcher's "Algebraic Topology."

1. Compute the simplicial homology groups $H_n(\mathbb{R}P^2; \mathbb{Z})$ using the Δ -complex structure given in class.
2. Hatcher, Exercise 4 on page 131.
3. Suppose $f: A \rightarrow B$ and $g: B \rightarrow C$ are homomorphisms of abelian groups. Show that there is an exact sequence

$$0 \rightarrow \ker(f) \rightarrow \ker(gf) \rightarrow \ker(g) \rightarrow \operatorname{coker}(f) \rightarrow \operatorname{coker}(gf) \rightarrow \operatorname{coker}(g) \rightarrow 0$$

4. In class, we defined subdivision maps $s_n^i: \Delta[n+1] \rightarrow \Delta[n] \times [0,1]$ for $0 \leq i \leq n$ by

$$s_n^i(t_1, \dots, t_{n+1}) = ((t_1, \dots, \widehat{t_{i+1}}, \dots, t_{n+1}), t_{i+1})$$

Show that these satisfy the relations

- $s_n^i d_{n+1}^j = \begin{cases} (d_n^{j-1}, \operatorname{id}) \circ s_{n-1}^i & \text{if } i < j-1 \\ (d_n^j, \operatorname{id}) \circ s_{n-1}^{i-1} & \text{if } i > j \end{cases}$
- $s_n^0 d_{n+1}^0 = i_0$
- $s_n^n d_{n+1}^{n+1} = i_1$
- $s_n^{i-1} d_{n+1}^i = s_n^i d_{n+1}^i$ for $1 \leq i < n+1$

Use this to show that the operator $h: C_n(X) \rightarrow C_{n+1}(X \times [0,1])$ given by

$$h\left(\sum a_\sigma \sigma\right) = \sum a_\sigma \sum_{i=0}^n (-1)^i (\sigma, \operatorname{id}) \circ s_n^i$$

satisfies $\partial h(x) + h \partial(x) = \tilde{i}_0(x) - \tilde{i}_1(x)$, where $\tilde{i}_k = (\sigma, \operatorname{id}) \circ i_k$.