

# Algebraic topology and arithmetic

Yifei Zhu

Southern University of Science and Technology

2019 Young Mathematician Forum, BICMR

# Motivation: cohomology theories and their operations

Generalized cohomology theory  $\{h^n\}: \text{Spaces} \rightarrow \text{AbGroups}$

Cup product  $\smile: h^*(X)$  a graded commutative algebra over  $h^*(\text{pt})$

Cohomology operation  $Q^i: h^*(-) \rightarrow h^{*+i}(-)$

Example (ordinary cohomology with  $\mathbb{Z}/2$ -coefficients)

Steenrod squares  $\text{Sq}^i: H^*(-; \mathbb{Z}/2) \rightarrow H^{*+i}(-; \mathbb{Z}/2)$

Power operation  $\text{Sq}^i(x) = x^2$  if  $i = |x|$

Steenrod algebra

# Motivation: cohomology theories and their operations

Generalized cohomology theory  $\{h^n\}: \text{Spaces} \rightarrow \text{AbGroups}$

Cup product  $\smile: h^*(X)$  a graded commutative algebra over  $h^*(\text{pt})$

Cohomology operation  $Q^i: h^*(-) \rightarrow h^{*+i}(-)$

Example (ordinary cohomology with  $\mathbb{Z}/2$ -coefficients)

Steenrod squares  $\text{Sq}^i: H^*(-; \mathbb{Z}/2) \rightarrow H^{*+i}(-; \mathbb{Z}/2)$

Power operation  $\text{Sq}^i(x) = x^2$  if  $i = |x|$

Steenrod algebra

# Motivation: cohomology theories and their operations

Generalized cohomology theory  $\{h^n\}: \text{Spaces} \rightarrow \text{AbGroups}$

Cup product  $\smile: h^*(X)$  a graded commutative algebra over  $h^*(\text{pt})$

Cohomology operation  $Q^i: h^*(-) \rightarrow h^{*+i}(-)$

Example (ordinary cohomology with  $\mathbb{Z}/2$ -coefficients)

Steenrod squares  $\text{Sq}^i: H^*(-; \mathbb{Z}/2) \rightarrow H^{*+i}(-; \mathbb{Z}/2)$

Power operation  $\text{Sq}^i(x) = x^2$  if  $i = |x|$

Steenrod algebra

# Motivation: cohomology theories and their operations

Generalized cohomology theory  $\{h^n\}: \text{Spaces} \rightarrow \text{AbGroups}$

Cup product  $\smile: h^*(X)$  a graded commutative algebra over  $h^*(\text{pt})$

Cohomology operation  $Q^i: h^*(-) \rightarrow h^{*+i}(-)$

Example (ordinary cohomology with  $\mathbb{Z}/2$ -coefficients)

Steenrod squares  $\text{Sq}^i: H^*(-; \mathbb{Z}/2) \rightarrow H^{*+i}(-; \mathbb{Z}/2)$

Power operation  $\text{Sq}^i(x) = x^2$  if  $i = |x|$

Steenrod algebra

# Motivation: cohomology theories and their operations

Generalized cohomology theory  $\{h^n\}: \text{Spaces} \rightarrow \text{AbGroups}$

Cup product  $\smile: h^*(X)$  a graded commutative algebra over  $h^*(\text{pt})$

Cohomology operation  $Q^i: h^*(-) \rightarrow h^{*+i}(-)$

Example (ordinary cohomology with  $\mathbb{Z}/2$ -coefficients)

Steenrod squares  $\text{Sq}^i: H^*(-; \mathbb{Z}/2) \rightarrow H^{*+i}(-; \mathbb{Z}/2)$

**Power operation**  $\text{Sq}^i(x) = x^2$  if  $i = |x|$

Steenrod algebra

# Motivation: cohomology theories and their operations

Generalized cohomology theory  $\{h^n\}: \text{Spaces} \rightarrow \text{AbGroups}$

Cup product  $\smile: h^*(X)$  a graded commutative algebra over  $h^*(\text{pt})$

Cohomology operation  $Q^i: h^*(-) \rightarrow h^{*+i}(-)$

Example (ordinary cohomology with  $\mathbb{Z}/2$ -coefficients)

Steenrod squares  $\text{Sq}^i: H^*(-; \mathbb{Z}/2) \rightarrow H^{*+i}(-; \mathbb{Z}/2)$

Power operation  $\text{Sq}^i(x) = x^2$  if  $i = |x|$

Steenrod algebra

# Motivation: cohomology theories and their operations

Generalized cohomology theory  $\{h^n\}: \text{Spaces} \rightarrow \text{AbGroups}$

Cup product  $\smile: h^*(X)$  a graded commutative algebra over  $h^*(\text{pt})$

Cohomology operation  $Q^i: h^*(-) \rightarrow h^{*+i}(-)$

## Example (ordinary cohomology with $\mathbb{Z}/2$ -coefficients)

Steenrod squares  $\text{Sq}^i: H^*(-; \mathbb{Z}/2) \rightarrow H^{*+i}(-; \mathbb{Z}/2)$

Power operation  $\text{Sq}^i(x) = x^2$  if  $i = |x|$

Adem relations  $\text{Sq}^i \text{Sq}^j = \sum_{k=0}^{\lfloor \frac{i}{2} \rfloor} \binom{j-k-1}{i-2k} \text{Sq}^{i+j-k} \text{Sq}^k, 0 < i < 2j$

Cartan formula  $\text{Sq}^i(xy) = \sum_{k=0}^i \text{Sq}^{i-k}(x) \text{Sq}^k(y)$



# Motivation: cohomology theories and their operations

Generalized cohomology theory  $\{h^n\}: \text{Spaces} \rightarrow \text{AbGroups}$

Cup product  $\smile: h^*(X)$  a graded commutative algebra over  $h^*(\text{pt})$

Cohomology operation  $Q^i: h^*(-) \rightarrow h^{*+i}(-)$

## Example (ordinary cohomology with $\mathbb{Z}/2$ -coefficients)

Steenrod squares  $\text{Sq}^i: H^*(-; \mathbb{Z}/2) \rightarrow H^{*+i}(-; \mathbb{Z}/2)$

Power operation  $\text{Sq}^i(x) = x^2$  if  $i = |x|$

Adem relations  $\text{Sq}^i \text{Sq}^j = \sum_{k=0}^{\lfloor \frac{i}{2} \rfloor} \binom{j-k-1}{i-2k} \text{Sq}^{i+j-k} \text{Sq}^k, 0 < i < 2j$

Cartan formula  $\text{Sq}^i(xy) = \sum_{k=0}^i \text{Sq}^{i-k}(x) \text{Sq}^k(y)$

# Motivation: cohomology theories and their operations

Generalized cohomology theory  $\{h^n\}: \text{Spaces} \rightarrow \text{AbGroups}$

Cup product  $\smile: h^*(X)$  a graded commutative algebra over  $h^*(\text{pt})$

Cohomology operation  $Q^i: h^*(-) \rightarrow h^{*+i}(-)$

## Example (ordinary cohomology with $\mathbb{Z}/2$ -coefficients)

Steenrod squares  $\text{Sq}^i: H^*(-; \mathbb{Z}/2) \rightarrow H^{*+i}(-; \mathbb{Z}/2)$

Power operation  $\text{Sq}^i(x) = x^2$  if  $i = |x|$

Adem relations  $\text{Sq}^i \text{Sq}^j = \sum_{k=0}^{\lfloor \frac{i}{2} \rfloor} \binom{j-k-1}{i-2k} \text{Sq}^{i+j-k} \text{Sq}^k, 0 < i < 2j$

Cartan formula  $\text{Sq}^i(xy) = \sum_{k=0}^i \text{Sq}^{i-k}(x) \text{Sq}^k(y)$

# Motivation: cohomology theories and their operations

Generalized cohomology theory  $\{h^n\}: \text{Spaces} \rightarrow \text{AbGroups}$

Cup product  $\smile: h^*(X)$  a graded commutative algebra over  $h^*(\text{pt})$

Cohomology operation  $Q^i: h^*(-) \rightarrow h^{*+i}(-)$

## Example (complex K-theory)

Adams operations  $\psi^i: K(-) \rightarrow K(-)$

Power operation  $\psi^p(x) \equiv x^p \pmod{p}$

$$\psi^i \psi^j = \psi^{ij}$$

$$\psi^i(xy) = \psi^i(x)\psi^i(y)$$

J. F. Adams, *Vector fields on spheres*, Ann. of Math. (2) **75** (1962)

# Motivation: cohomology theories and their operations

Generalized cohomology theory  $\{h^n\}: \text{Spaces} \rightarrow \text{AbGroups}$

Cup product  $\rightsquigarrow h^*(X)$  a graded commutative algebra over  $h^*(\text{pt})$

Cohomology operation  $Q^i: h^*(-) \rightarrow h^{*+i}(-)$

## Example (more – a sample)

Voevodsky, *Reduced power operations in motivic cohomology*, 2003.

Lipshitz and Sarkar, *A Steenrod square on Khovanov homology*, 2014.

Feng, *Étale Steenrod operations and the Artin–Tate pairing*, 2018.

Seidel, *Formal groups and quantum cohomology*, 2019.

# Background: Chromatic Homotopy Theory 色展同伦论

## A connection between Topology and Arithmetic (Quillen '69)

stable homotopy theory  $\longleftrightarrow$  1-dim formal group laws

complex-oriented  $h^*(-)$   $F(x, y)$  over  $h^*(\text{pt})$

$$c_1(L_1 \otimes L_2) = F(c_1(L_1), c_1(L_2))$$

### Example

$$H^*(-; \mathbb{Z}) \longleftrightarrow \mathbb{G}_a(x, y) = x + y$$

$$K^*(-) \longleftrightarrow \mathbb{G}_m(x, y) = x + y - xy = 1 - (1 - x)(1 - y)$$

# Background: Chromatic Homotopy Theory 色展同伦论

## A connection between Topology and Arithmetic (Quillen '69)

stable homotopy theory  $\longleftrightarrow$  1-dim formal group laws

complex-oriented  $h^*(-)$   $F(x, y)$  over  $h^*(\text{pt})$

$$c_1(L_1 \otimes L_2) = F(c_1(L_1), c_1(L_2))$$

### Example

$$H^*(-; \mathbb{Z}) \longleftrightarrow \mathbb{G}_a(x, y) = x + y$$

$$K^*(-) \longleftrightarrow \mathbb{G}_m(x, y) = x + y - xy = 1 - (1 - x)(1 - y)$$

# Background: Chromatic Homotopy Theory 色展同伦论

## A connection between Topology and Arithmetic (Quillen '69)

stable homotopy theory  $\longleftrightarrow$  1-dim formal group laws

complex-oriented  $h^*(-)$   $F(x, y)$  over  $h^*(\text{pt})$

$$c_1(L_1 \otimes L_2) = F(c_1(L_1), c_1(L_2))$$

### Example

$$H^*(-; \mathbb{Z}) \longleftrightarrow \mathbb{G}_a(x, y) = x + y$$

$$K^*(-) \longleftrightarrow \mathbb{G}_m(x, y) = x + y - xy = 1 - (1 - x)(1 - y)$$

# Background: Chromatic Homotopy Theory 色展同伦论

## A connection between Topology and Arithmetic (Quillen '69)

stable homotopy theory  $\longleftrightarrow$  1-dim formal group laws

complex-oriented  $h^*(-)$   $F(x, y)$  over  $h^*(\text{pt})$

$$c_1(L_1 \otimes L_2) = F(c_1(L_1), c_1(L_2))$$

### Example

$$H^*(-; \mathbb{Z}) \longleftrightarrow \mathbb{G}_a(x, y) = x + y$$

$$K^*(-) \longleftrightarrow \mathbb{G}_m(x, y) = x + y - xy = 1 - (1 - x)(1 - y)$$



# Background: Chromatic Homotopy Theory 色展同伦论

## A connection between Topology and Arithmetic (Quillen '69)

stable homotopy theory  $\longleftrightarrow$  1-dim formal group laws

complex-oriented  $h^*(-)$   $F(x, y)$  over  $h^*(\text{pt})$

$$c_1(L_1 \otimes L_2) = F(c_1(L_1), c_1(L_2))$$

### Example

$$H^*(-; \mathbb{Z}) \longleftrightarrow \mathbb{G}_a(x, y) = x + y$$

$$K^*(-) \longleftrightarrow \mathbb{G}_m(x, y) = x + y - xy = 1 - (1 - x)(1 - y)$$

# Background: Chromatic Homotopy Theory 色展同伦论

## A connection between Topology and Arithmetic (Quillen '69)

stable homotopy theory  $\longleftrightarrow$  1-dim formal group laws

complex-oriented  $h^*(-)$   $F(x, y)$  over  $h^*(\text{pt})$

$$c_1(L_1 \otimes L_2) = F(c_1(L_1), c_1(L_2))$$

### Example

$$H^*(-; \mathbb{Z}) \longleftrightarrow \mathbb{G}_a(x, y) = x + y$$

$$K^*(-) \longleftrightarrow \mathbb{G}_m(x, y) = x + y - xy = 1 - (1 - x)(1 - y)$$

# Background: Chromatic Homotopy Theory 色展同伦论

## A connection between Topology and Arithmetic (Quillen '69)

stable homotopy theory  $\longleftrightarrow$  1-dim formal group laws

complex-oriented  $h^*(-)$   $F(x, y)$  over  $h^*(\text{pt})$

$$c_1(L_1 \otimes L_2) = F(c_1(L_1), c_1(L_2))$$

### Example

$$H^*(-; \mathbb{Z}) \longleftrightarrow \mathbb{G}_a(x, y) = x + y$$

$$K^*(-) \longleftrightarrow \mathbb{G}_m(x, y) = x + y - xy = 1 - (1 - x)(1 - y)$$

# Background: Chromatic Homotopy Theory 色展同伦论

## A connection between Topology and Arithmetic (Quillen '69)

stable homotopy theory  $\longleftrightarrow$  1-dim formal group laws

complex-oriented  $h^*(-)$   $F(x, y)$  over  $h^*(\text{pt})$

$$c_1(L_1 \otimes L_2) = F(c_1(L_1), c_1(L_2))$$

### Example

$$H^*(-; \mathbb{Z}) \longleftrightarrow \mathbb{G}_a(x, y) = x + y$$

$$K^*(-) \longleftrightarrow \mathbb{G}_m(x, y) = x + y - xy = 1 - (1 - x)(1 - y)$$

# Elliptic cohomology and Morava E-theory

Definition (Ando–Hopkins–Strickland '01, Lurie '09, '18)

$$\text{elliptic cohomology theory} = \left\{ \begin{array}{l} E, \quad C_{E^0(\text{pt})}, \\ \alpha: \text{Spf } E^0(\mathbb{CP}^\infty) \xrightarrow{\sim} \widehat{C} \end{array} \right\}$$

Theorem (Morava '78, Goerss–Hopkins–Miller '90s–'04)

$\mathcal{E}: \{\text{formal groups over perfect fields, isos}\} \rightarrow \{\mathbb{E}_\infty\text{-ring spectra}\}$

- $\text{Spf } E^0(\mathbb{CP}^\infty) =$  the univ deformation of a fg  $F$  of height  $n$  over a perfect field  $k$  of char  $p$
- $\pi_* E \cong \mathbb{W}(k)[[u_1, \dots, u_{n-1}]] [u^{\pm 1}], \quad |u| = -2$

# Elliptic cohomology and Morava E-theory

Definition (Ando–Hopkins–Strickland '01, Lurie '09, '18)

$$\text{elliptic cohomology theory} = \left\{ \begin{array}{l} E, \quad C_{E^0(\text{pt})}, \\ \alpha: \text{Spf } E^0(\mathbb{CP}^\infty) \xrightarrow{\sim} \widehat{C} \end{array} \right\}$$

Theorem (Morava '78, Goerss–Hopkins–Miller '90s–'04)

$\mathcal{E}: \{\text{formal groups over perfect fields, isos}\} \rightarrow \{\mathbb{E}_\infty\text{-ring spectra}\}$

- $\text{Spf } E^0(\mathbb{CP}^\infty) =$  the univ deformation of a fg  $F$  of height  $n$  over a perfect field  $k$  of char  $p$
- $\pi_* E \cong \mathbb{W}(k)[[u_1, \dots, u_{n-1}]] [u^{\pm 1}], \quad |u| = -2$

# Elliptic cohomology and Morava E-theory

Definition (Ando–Hopkins–Strickland '01, Lurie '09, '18)

$$\text{elliptic cohomology theory} = \left\{ \begin{array}{l} E, \quad C_{E^0(\text{pt})}, \\ \alpha: \text{Spf } E^0(\mathbb{CP}^\infty) \xrightarrow{\sim} \widehat{C} \end{array} \right\}$$

Theorem (Morava '78, Goerss–Hopkins–Miller '90s–'04)

$\mathcal{E}: \{\text{formal groups over perfect fields, isos}\} \rightarrow \{\mathbb{E}_\infty\text{-ring spectra}\}$

- $\text{Spf } E^0(\mathbb{CP}^\infty) =$  the univ deformation of a fg  $F$  of height  $n$  over a perfect field  $k$  of char  $p$
- $\pi_* E \cong \mathbb{W}(k)[[u_1, \dots, u_{n-1}]] [u^{\pm 1}], \quad |u| = -2$

# Elliptic cohomology and Morava E-theory

Definition (Ando–Hopkins–Strickland '01, Lurie '09, '18)

$$\text{elliptic cohomology theory} = \left\{ \begin{array}{l} E, \quad C_{E^0(\text{pt})}, \\ \alpha: \text{Spf } E^0(\mathbb{CP}^\infty) \xrightarrow{\sim} \widehat{C} \end{array} \right\}$$

Theorem (Morava '78, Goerss–Hopkins–Miller '90s–'04)

$\mathcal{E}: \{\text{formal groups over perfect fields, isos}\} \rightarrow \{\mathbb{E}_\infty\text{-ring spectra}\}$

- $\text{Spf } E^0(\mathbb{CP}^\infty) =$  the univ deformation of a fg  $F$  of height  $n$  over a perfect field  $k$  of char  $p$
- $\pi_* E \cong \mathbb{W}(k)[[u_1, \dots, u_{n-1}]] [u^{\pm 1}], \quad |u| = -2$



# Elliptic cohomology and Morava E-theory

Definition (Ando–Hopkins–Strickland '01, Lurie '09, '18)

$$\text{elliptic cohomology theory} = \left\{ \begin{array}{l} E, \quad C_{E^0(\text{pt})}, \\ \alpha: \text{Spf } E^0(\mathbb{CP}^\infty) \xrightarrow{\sim} \widehat{C} \end{array} \right\}$$

Theorem (Morava '78, Goerss–Hopkins–Miller '90s–'04)

$\mathcal{E}: \{\text{formal groups over perfect fields, isos}\} \rightarrow \{\mathbb{E}_\infty\text{-ring spectra}\}$

- $\text{Spf } E^0(\mathbb{CP}^\infty) =$  the univ deformation of a fg  $F$  of height  $n$  over a perfect field  $k$  of char  $p$
- $\pi_* E \cong \mathbb{W}(k)[[u_1, \dots, u_{n-1}]] [u^{\pm 1}], \quad |u| = -2$

# Elliptic cohomology and Morava E-theory

Definition (Ando–Hopkins–Strickland '01, Lurie '09, '18)

$$\text{elliptic cohomology theory} = \left\{ \begin{array}{l} E, \quad C_{E^0(\text{pt})}, \\ \alpha: \text{Spf } E^0(\mathbb{CP}^\infty) \xrightarrow{\sim} \widehat{C} \end{array} \right\}$$

Theorem (Morava '78, Goerss–Hopkins–Miller '90s–'04)

$\mathcal{E}: \{\text{formal groups over perfect fields, isos}\} \rightarrow \{\mathbb{E}_\infty\text{-ring spectra}\}$

- $\text{Spf } E^0(\mathbb{CP}^\infty) =$  the univ deformation of a fg  $F$  of height  $n$  over a perfect field  $k$  of char  $p$
- $\pi_* E \cong \mathbb{W}(k)[[u_1, \dots, u_{n-1}]] [u^{\pm 1}], \quad |u| = -2$

# Elliptic cohomology and Morava E-theory

Definition (Ando–Hopkins–Strickland '01, Lurie '09, '18)

$$\text{elliptic cohomology theory} = \left\{ \begin{array}{l} E, \quad C_{E^0(\text{pt})}, \\ \alpha: \text{Spf } E^0(\mathbb{CP}^\infty) \xrightarrow{\sim} \widehat{C} \end{array} \right\}$$

Theorem (Morava '78, Goerss–Hopkins–Miller '90s–'04)

$\mathcal{E}: \{\text{formal groups over perfect fields, isos}\} \rightarrow \{\mathbb{E}_\infty\text{-ring spectra}\}$

- $\text{Spf } E^0(\mathbb{CP}^\infty) =$  the univ deformation of a fg  $F$  of height  $n$  over a perfect field  $k$  of char  $p$
- $\pi_* E \cong \mathbb{W}(k)[[u_1, \dots, u_{n-1}]] [u^{\pm 1}], \quad |u| = -2$

# Elliptic cohomology and Morava E-theory

Definition (Ando–Hopkins–Strickland '01, Lurie '09, '18)

$$\text{elliptic cohomology theory} = \left\{ \begin{array}{l} E, \quad C_{E^0(\text{pt})}, \\ \alpha: \text{Spf } E^0(\mathbb{CP}^\infty) \xrightarrow{\sim} \widehat{C} \end{array} \right\}$$

Theorem (Morava '78, Goerss–Hopkins–Miller '90s–'04)

$\mathcal{E}: \{\text{formal groups over perfect fields, isos}\} \rightarrow \{\mathbb{E}_\infty\text{-ring spectra}\}$

- $\text{Spf } E^0(\mathbb{CP}^\infty) =$  the univ deformation of a fg  $F$  of height  $n$  over a perfect field  $k$  of char  $p$
- $\pi_* E \cong \mathbb{W}(k)[[u_1, \dots, u_{n-1}]] [u^{\pm 1}], \quad |u| = -2$

# Elliptic cohomology and Morava E-theory

Definition (Ando–Hopkins–Strickland '01, Lurie '09, '18)

$$\text{elliptic cohomology theory} = \left\{ \begin{array}{l} E, \quad C_{E^0(\text{pt})}, \\ \alpha: \text{Spf } E^0(\mathbb{CP}^\infty) \xrightarrow{\sim} \widehat{C} \end{array} \right\}$$

Theorem (Morava '78, Goerss–Hopkins–Miller '90s–'04)

$\mathcal{E}: \{\text{formal groups over perfect fields, isos}\} \rightarrow \{\mathbb{E}_\infty\text{-ring spectra}\}$

- $\text{Spf } E^0(\mathbb{CP}^\infty) =$  the univ deformation of a fg  $F$  of height  $n$  over a perfect field  $k$  of char  $p$
- $\pi_* E \cong \mathbb{W}(k)[[u_1, \dots, u_{n-1}]] [u^{\pm 1}], \quad |u| = -2$

# Elliptic cohomology and Morava E-theory

Definition (Ando–Hopkins–Strickland '01, Lurie '09, '18)

$$\text{elliptic cohomology theory} = \left\{ \begin{array}{l} E, \quad C_{E^0(\text{pt})}, \\ \alpha: \text{Spf } E^0(\mathbb{CP}^\infty) \xrightarrow{\sim} \widehat{C} \end{array} \right\}$$

Theorem (Morava '78, Goerss–Hopkins–Miller '90s–'04)

$\mathcal{E}: \{\text{formal groups over perfect fields, isos}\} \rightarrow \{\mathbb{E}_\infty\text{-ring spectra}\}$

- $\text{Spf } E^0(\mathbb{CP}^\infty) =$  the univ deformation of a fg  $F$  of height  $n$  over a perfect field  $k$  of char  $p$
- $\pi_* E \cong \mathbb{W}(k)[[u_1, \dots, u_{n-1}]] [u^{\pm 1}], \quad |u| = -2$

# Elliptic cohomology and Morava E-theory

Definition (Ando–Hopkins–Strickland '01, Lurie '09, '18)

$$\text{elliptic cohomology theory} = \left\{ \begin{array}{l} E, \quad C_{E^0(\text{pt})}, \\ \alpha: \text{Spf } E^0(\mathbb{CP}^\infty) \xrightarrow{\sim} \widehat{C} \end{array} \right\}$$

Theorem (Morava '78, Goerss–Hopkins–Miller '90s–'04)

$\mathcal{E}: \{\text{formal groups over perfect fields, isos}\} \rightarrow \{\mathbb{E}_\infty\text{-ring spectra}\}$

- $\text{Spf } E^0(\mathbb{CP}^\infty) = \text{the univ deformation}$  of a fg  $F$  of height  $n$  over a perfect field  $k$  of char  $p$
- $\pi_* E \cong \mathbb{W}(k)[[u_1, \dots, u_{n-1}]] [u^{\pm 1}]$ ,  $|u| = -2$

# Elliptic cohomology and Morava E-theory

Definition (Ando–Hopkins–Strickland '01, Lurie '09, '18)

$$\text{elliptic cohomology theory} = \left\{ \begin{array}{l} E, \quad C_{E^0(\text{pt})}, \\ \alpha: \text{Spf } E^0(\mathbb{CP}^\infty) \xrightarrow{\sim} \widehat{C} \end{array} \right\}$$

Theorem (Morava '78, Goerss–Hopkins–Miller '90s–'04)

$$\mathcal{E}: \{\text{formal groups over perfect fields, isos}\} \rightarrow \{\mathbb{E}_\infty\text{-ring spectra}\}$$

$E$  = Morava E-theory of height  $n$  at the prime  $p$

Goal explore the structure on  $E^*(-)$ .      Topology  $\longleftrightarrow$  Arithmetic



# Elliptic cohomology and Morava E-theory

Definition (Ando–Hopkins–Strickland '01, Lurie '09, '18)

$$\text{elliptic cohomology theory} = \left\{ \begin{array}{l} E, \quad C_{E^0(\text{pt})}, \\ \alpha: \text{Spf } E^0(\mathbb{CP}^\infty) \xrightarrow{\sim} \widehat{C} \end{array} \right\}$$

Theorem (Morava '78, Goerss–Hopkins–Miller '90s–'04)

$$\mathcal{E}: \{\text{formal groups over perfect fields, isos}\} \rightarrow \{\mathbb{E}_\infty\text{-ring spectra}\}$$

$E$  = Morava E-theory of height  $n$  at the prime  $p$

Goal explore the structure on  $E^*(-)$ .      **Topology**  $\longleftrightarrow$  **Arithmetic**

# Elliptic cohomology and Morava E-theory

Definition (Ando–Hopkins–Strickland '01, Lurie '09, '18)

$$\text{elliptic cohomology theory} = \left\{ \begin{array}{l} E, \quad C_{E^0(\text{pt})}, \\ \alpha: \text{Spf } E^0(\mathbb{CP}^\infty) \xrightarrow{\sim} \widehat{C} \end{array} \right\}$$

Theorem (Morava '78, Goerss–Hopkins–Miller '90s–'04)

$$\mathcal{E}: \{\text{formal groups over perfect fields, isos}\} \rightarrow \{\mathbb{E}_\infty\text{-ring spectra}\}$$

$E$  = Morava E-theory of height  $n$  at the prime  $p$

Goal explore the structure on  $E^*(-)$ . **Topology**  $\longleftrightarrow$  **Arithmetic**

# Power operations for Morava E-theory

$$M = E\text{-module} \quad \pi_0 M = [S, M]_S \cong [E, M]_E$$

$$\mathbb{P}_E(M) = \bigvee_{i \geq 0} \mathbb{P}_E^i(M) = \bigvee_{i \geq 0} \underbrace{(M \wedge_E \cdots \wedge_E M)}_{i\text{-fold}}_{h\Sigma_i}$$

$A =$  commutative  $E$ -algebra

$=$  algebra for the monad  $\mathbb{P}_E$  with  $\mu: \mathbb{P}_E(A) \rightarrow A$

total power operation  $\psi^i: \pi_0 A \rightarrow \pi_0(A^{B\Sigma_i^+})$   
 $\forall \eta \in \pi_0 \mathbb{P}_E^i(E)$ , individual po  $Q_\eta: \pi_0 A \rightarrow \pi_0 A$   $\left. \vphantom{\begin{matrix} \psi^i \\ Q_\eta \end{matrix}} \right\} \xrightarrow{/I} \text{additive}$

$$E \xrightarrow{f_\eta} \mathbb{P}_E^i(E) \xrightarrow{\mathbb{P}_E^i(f_x)} \mathbb{P}_E^i(A) \hookrightarrow \mathbb{P}_E(A) \xrightarrow{\mu} A$$

# Power operations for Morava E-theory

$$M = E\text{-module} \quad \pi_0 M = [S, M]_S \cong [E, M]_E$$

$$\mathbb{P}_E(M) = \bigvee_{i \geq 0} \mathbb{P}_E^i(M) = \bigvee_{i \geq 0} \underbrace{(M \wedge_E \cdots \wedge_E M)_{h\Sigma_i}}_{i\text{-fold}}$$

$A =$  commutative  $E$ -algebra

$=$  algebra for the monad  $\mathbb{P}_E$  with  $\mu: \mathbb{P}_E(A) \rightarrow A$

total power operation  $\psi^i: \pi_0 A \rightarrow \pi_0(A^{B\Sigma_i^+})$  }  $\xrightarrow{/I}$  additive  
 $\forall \eta \in \pi_0 \mathbb{P}_E^i(E), \text{ individual po } Q_\eta: \pi_0 A \rightarrow \pi_0 A$

$$E \xrightarrow{f_\eta} \mathbb{P}_E^i(E) \xrightarrow{\mathbb{P}_E^i(f_x)} \mathbb{P}_E^i(A) \hookrightarrow \mathbb{P}_E(A) \xrightarrow{\mu} A$$

# Power operations for Morava E-theory

$$M = E\text{-module} \quad \pi_0 M = [S, M]_S \cong [E, M]_E$$

$$\mathbb{P}_E(M) = \bigvee_{i \geq 0} \mathbb{P}_E^i(M) = \bigvee_{i \geq 0} \underbrace{(M \wedge_E \cdots \wedge_E M)_{h\Sigma_i}}_{i\text{-fold}}$$

$A =$  commutative  $E$ -algebra

$=$  algebra for the monad  $\mathbb{P}_E$  with  $\mu: \mathbb{P}_E(A) \rightarrow A$

total power operation  $\psi^i: \pi_0 A \rightarrow \pi_0(A^{B\Sigma_i^+})$   $\left. \vphantom{\begin{matrix} \psi^i \\ \forall \eta \in \pi_0 \mathbb{P}_E^i(E), \text{ individual po } Q_\eta: \pi_0 A \rightarrow \pi_0 A \end{matrix}} \right\} \xrightarrow{/I} \text{additive}$

$$E \xrightarrow{f_\eta} \mathbb{P}_E^i(E) \xrightarrow{\mathbb{P}_E^i(f_x)} \mathbb{P}_E^i(A) \hookrightarrow \mathbb{P}_E(A) \xrightarrow{\mu} A$$

# Power operations for Morava E-theory

$$M = E\text{-module} \quad \pi_0 M = [S, M]_S \cong [E, M]_E$$

$$\mathbb{P}_E(M) = \bigvee_{i \geq 0} \mathbb{P}_E^i(M) = \bigvee_{i \geq 0} \underbrace{(M \wedge_E \cdots \wedge_E M)_{h\Sigma_i}}_{i\text{-fold}}$$

$A =$  commutative  $E$ -algebra

$=$  algebra for the monad  $\mathbb{P}_E$  with  $\mu: \mathbb{P}_E(A) \rightarrow A$

total power operation  $\psi^i: \pi_0 A \rightarrow \pi_0(A^{B\Sigma_i^+})$  }  $\xrightarrow{/I}$  additive  
 $\forall \eta \in \pi_0 \mathbb{P}_E^i(E), \text{ individual po } Q_\eta: \pi_0 A \rightarrow \pi_0 A$

$$E \xrightarrow{f_\eta} \mathbb{P}_E^i(E) \xrightarrow{\mathbb{P}_E^i(f_x)} \mathbb{P}_E^i(A) \hookrightarrow \mathbb{P}_E(A) \xrightarrow{\mu} A$$

# Power operations for Morava E-theory

$$M = E\text{-module} \quad \pi_0 M = [S, M]_S \cong [E, M]_E$$

$$\mathbb{P}_E(M) = \bigvee_{i \geq 0} \mathbb{P}_E^i(M) = \bigvee_{i \geq 0} \underbrace{(M \wedge_E \cdots \wedge_E M)_{h\Sigma_i}}_{i\text{-fold}}$$

$A =$  commutative  $E$ -algebra

$=$  algebra for the monad  $\mathbb{P}_E$  with  $\mu: \mathbb{P}_E(A) \rightarrow A$

total power operation  $\psi^i: \pi_0 A \rightarrow \pi_0(A^{B\Sigma_i^+})$   $\left. \vphantom{\begin{matrix} \psi^i \\ \forall \eta \in \pi_0 \mathbb{P}_E^i(E), \text{ individual po } Q_\eta: \pi_0 A \rightarrow \pi_0 A \end{matrix}} \right\} \xrightarrow{/I} \text{additive}$

$$E \xrightarrow{f_\eta} \mathbb{P}_E^i(E) \xrightarrow{\mathbb{P}_E^i(f_x)} \mathbb{P}_E^i(A) \hookrightarrow \mathbb{P}_E(A) \xrightarrow{\mu} A$$

# Power operations for Morava E-theory

$$M = E\text{-module} \quad \pi_0 M = [S, M]_S \cong [E, M]_E$$

$$\mathbb{P}_E(M) = \bigvee_{i \geq 0} \mathbb{P}_E^i(M) = \bigvee_{i \geq 0} \underbrace{(M \wedge_E \cdots \wedge_E M)_{h\Sigma_i}}_{i\text{-fold}}$$

$A =$  commutative  $E$ -algebra

$=$  algebra for the monad  $\mathbb{P}_E$  with  $\mu: \mathbb{P}_E(A) \rightarrow A$

total power operation  $\psi^i: \pi_0 A \rightarrow \pi_0(A^{B\Sigma_i^+})$   
 $\forall \eta \in \pi_0 \mathbb{P}_E^i(E)$ , individual po  $Q_\eta: \pi_0 A \rightarrow \pi_0 A$   $\left. \vphantom{\begin{matrix} \psi^i \\ Q_\eta \end{matrix}} \right\} \xrightarrow{/I} \text{additive}$

$$E \xrightarrow{f_\eta} \mathbb{P}_E^i(E) \xrightarrow{\mathbb{P}_E^i(f_x)} \mathbb{P}_E^i(A) \hookrightarrow \mathbb{P}_E(A) \xrightarrow{\mu} A$$



# Power operations for Morava E-theory

$$M = E\text{-module} \quad \pi_0 M = [S, M]_S \cong [E, M]_E$$

$$\mathbb{P}_E(M) = \bigvee_{i \geq 0} \mathbb{P}_E^i(M) = \bigvee_{i \geq 0} \underbrace{(M \wedge_E \cdots \wedge_E M)_{h\Sigma_i}}_{i\text{-fold}}$$

$A =$  commutative  $E$ -algebra

$=$  algebra for the monad  $\mathbb{P}_E$  with  $\mu: \mathbb{P}_E(A) \rightarrow A$

total power operation  $\psi^i: \pi_0 A \rightarrow \pi_0(A^{B\Sigma_i^+})$   
 $\forall \eta \in \pi_0 \mathbb{P}_E^i(E), \text{ individual po } Q_\eta: \pi_0 A \rightarrow \pi_0 A \quad \left. \vphantom{\begin{matrix} \psi^i \\ Q_\eta \end{matrix}} \right\} \xrightarrow{/I} \text{additive}$

$$E \xrightarrow{f_\eta} \mathbb{P}_E^i(E) \xrightarrow{\mathbb{P}_E^i(f_{\mathbf{x}})} \mathbb{P}_E^i(A) \hookrightarrow \mathbb{P}_E(A) \xrightarrow{\mu} A$$

# Power operations for Morava E-theory

$$M = E\text{-module} \quad \pi_0 M = [S, M]_S \cong [E, M]_E$$

$$\mathbb{P}_E(M) = \bigvee_{i \geq 0} \mathbb{P}_E^i(M) = \bigvee_{i \geq 0} \underbrace{(M \wedge_E \cdots \wedge_E M)_{h\Sigma_i}}_{i\text{-fold}}$$

$A =$  commutative  $E$ -algebra

$=$  algebra for the monad  $\mathbb{P}_E$  with  $\mu: \mathbb{P}_E(A) \rightarrow A$

total power operation  $\psi^i: \pi_0 A \rightarrow \pi_0(A^{B\Sigma_i^+})$   
 $\forall \eta \in \pi_0 \mathbb{P}_E^i(E)$ , individual po  $Q_\eta: \pi_0 A \rightarrow \pi_0 A$   $\left. \vphantom{\begin{matrix} \psi^i \\ Q_\eta \end{matrix}} \right\} \xrightarrow{/I} \text{additive}$

$$E \xrightarrow{f_\eta} \mathbb{P}_E^i(E) \xrightarrow{\mathbb{P}_E^i(f_x)} \mathbb{P}_E^i(A) \hookrightarrow \mathbb{P}_E(A) \xrightarrow{\mu} A$$

# Power operations for Morava E-theory

$$M = E\text{-module} \quad \pi_0 M = [S, M]_S \cong [E, M]_E$$

$$\mathbb{P}_E(M) = \bigvee_{i \geq 0} \mathbb{P}_E^i(M) = \bigvee_{i \geq 0} \underbrace{(M \wedge_E \cdots \wedge_E M)_{h\Sigma_i}}_{i\text{-fold}}$$

$A =$  commutative  $E$ -algebra

$=$  algebra for the monad  $\mathbb{P}_E$  with  $\mu: \mathbb{P}_E(A) \rightarrow A$

total power operation  $\psi^i: \pi_0 A \rightarrow \pi_0(A^{B\Sigma_i^+})$   
 $\forall \eta \in \pi_0 \mathbb{P}_E^i(E)$ , individual po  $Q_\eta: \pi_0 A \rightarrow \pi_0 A$   $\left. \vphantom{\begin{matrix} \psi^i \\ Q_\eta \end{matrix}} \right\} \xrightarrow{/I} \text{additive}$

$$E \xrightarrow{f_\eta} \mathbb{P}_E^i(E) \xrightarrow{\mathbb{P}_E^i(f_x)} \mathbb{P}_E^i(A) \hookrightarrow \mathbb{P}_E(A) \xrightarrow{\mu} A$$

# Power operations for Morava E-theory

$$M = E\text{-module} \quad \pi_0 M = [S, M]_S \cong [E, M]_E$$

$$\mathbb{P}_E(M) = \bigvee_{i \geq 0} \mathbb{P}_E^i(M) = \bigvee_{i \geq 0} \underbrace{(M \wedge_E \cdots \wedge_E M)_{h\Sigma_i}}_{i\text{-fold}}$$

$A =$  commutative  $E$ -algebra

$=$  algebra for the monad  $\mathbb{P}_E$  with  $\mu: \mathbb{P}_E(A) \rightarrow A$

total power operation  $\psi^i: \pi_0 A \rightarrow \pi_0(A^{B\Sigma_i^+})$   
 $\forall \eta \in \pi_0 \mathbb{P}_E^i(E)$ , individual po  $Q_\eta: \pi_0 A \rightarrow \pi_0 A$  }  $\xrightarrow{/I}$  additive

$$E \xrightarrow{f_\eta} \mathbb{P}_E^i(E) \xrightarrow{\mathbb{P}_E^i(f_x)} \mathbb{P}_E^i(A) \hookrightarrow \mathbb{P}_E(A) \xrightarrow{\mu} A$$

# Power operations for Morava E-theory

$$M = E\text{-module} \quad \pi_0 M = [S, M]_S \cong [E, M]_E$$

$$\mathbb{P}_E(M) = \bigvee_{i \geq 0} \mathbb{P}_E^i(M) = \bigvee_{i \geq 0} \underbrace{(M \wedge_E \cdots \wedge_E M)_{h\Sigma_i}}_{i\text{-fold}}$$

$A =$  commutative  $E$ -algebra

$=$  algebra for the monad  $\mathbb{P}_E$  with  $\mu: \mathbb{P}_E(A) \rightarrow A$

total power operation  $\psi^i: \pi_0 A \rightarrow \pi_0(A^{B\Sigma_i^+})$   
 $\forall \eta \in \pi_0 \mathbb{P}_E^i(E)$ , individual po  $Q_\eta: \pi_0 A \rightarrow \pi_0 A$   $\left. \vphantom{\begin{matrix} \psi^i \\ Q_\eta \end{matrix}} \right\} \xrightarrow{/I} \text{additive}$

$$E \xrightarrow{f_\eta} \mathbb{P}_E^i(E) \xrightarrow{\mathbb{P}_E^i(f_x)} \mathbb{P}_E^i(A) \hookrightarrow \mathbb{P}_E(A) \xrightarrow{\mu} A$$

# Power operations for Morava E-theory

$$M = E\text{-module} \quad \pi_0 M = [S, M]_S \cong [E, M]_E$$

$$\mathbb{P}_E(M) = \bigvee_{i \geq 0} \mathbb{P}_E^i(M) = \bigvee_{i \geq 0} \underbrace{(M \wedge_E \cdots \wedge_E M)_{h\Sigma_i}}_{i\text{-fold}}$$

$A =$  commutative  $E$ -algebra

$=$  algebra for the monad  $\mathbb{P}_E$  with  $\mu: \mathbb{P}_E(A) \rightarrow A$

total power operation  $\psi^i: \pi_0 A \rightarrow \pi_0(A^{B\Sigma_i^+})$   
 $\forall \eta \in \pi_0 \mathbb{P}_E^i(E)$ , individual po  $Q_\eta: \pi_0 A \rightarrow \pi_0 A$  }  $\xrightarrow{/I}$  additive

$$E \xrightarrow{f_\eta} \mathbb{P}_E^i(E) \xrightarrow{\mathbb{P}_E^i(f_x)} \mathbb{P}_E^i(A) \hookrightarrow \mathbb{P}_E(A) \xrightarrow{\mu} A$$

# Power operations for Morava E-theory

$$M = E\text{-module} \quad \pi_0 M = [S, M]_S \cong [E, M]_E$$

$$\mathbb{P}_E(M) = \bigvee_{i \geq 0} \mathbb{P}_E^i(M) = \bigvee_{i \geq 0} \underbrace{(M \wedge_E \cdots \wedge_E M)_{h\Sigma_i}}_{i\text{-fold}}$$

$A =$  commutative  $E$ -algebra

$=$  algebra for the monad  $\mathbb{P}_E$  with  $\mu: \mathbb{P}_E(A) \rightarrow A$

total power operation  $\psi^i: \pi_0 A \rightarrow \pi_0(A^{B\Sigma_i^+})$   
 $\forall \eta \in \pi_0 \mathbb{P}_E^i(E)$ , individual po  $Q_\eta: \pi_0 A \rightarrow \pi_0 A$   $\left. \vphantom{\begin{matrix} \psi^i \\ Q_\eta \end{matrix}} \right\} \xrightarrow{/I} \text{additive}$

$$E \xrightarrow{f_\eta} \mathbb{P}_E^i(E) \xrightarrow{\mathbb{P}_E^i(f_x)} \mathbb{P}_E^i(A) \hookrightarrow \mathbb{P}_E(A) \xrightarrow{\mu} A$$

# Power operations for Morava E-theory

$$M = E\text{-module} \quad \pi_0 M = [S, M]_S \cong [E, M]_E$$

$$\mathbb{P}_E(M) = \bigvee_{i \geq 0} \mathbb{P}_E^i(M) = \bigvee_{i \geq 0} \underbrace{(M \wedge_E \cdots \wedge_E M)_{h\Sigma_i}}_{i\text{-fold}}$$

$A =$  commutative  $E$ -algebra

$=$  algebra for the monad  $\mathbb{P}_E$  with  $\mu: \mathbb{P}_E(A) \rightarrow A$

total power operation  $\psi^i: \pi_0 A \rightarrow \pi_0(A^{B\Sigma_i^+})$   
 $\forall \eta \in \pi_0 \mathbb{P}_E^i(E), \text{ individual po } Q_\eta: \pi_0 A \rightarrow \pi_0 A \quad \left. \vphantom{\begin{matrix} \psi^i \\ Q_\eta \end{matrix}} \right\} \xrightarrow{/I} \text{additive}$

$$E \xrightarrow{f_\eta} \mathbb{P}_E^i(E) \xrightarrow{\mathbb{P}_E^i(f_x)} \mathbb{P}_E^i(A) \hookrightarrow \mathbb{P}_E(A) \xrightarrow{\mu} A$$



# Power operations for Morava E-theory (height $n$ prime $p$ )

Theorem (Rezk '09, Barthel–Frankland '13)

If  $A = K(n)$ -local commutative  $E$ -algebra, then

$\pi_* A =$  graded amplified  $L$ -complete  $\Gamma$ -ring

- $\Gamma =$  twisted bialgebra over  $E_0$  (Dyer–Lashof algebra)
- $\exists Q_0 \in \Gamma$  with  $Q_0(x) \equiv x^p \pmod{p}$  (Frobenius congruence)

Goal make this structure explicit just as for Dyer–Lashof/Steenrod operations in ordinary homology.

The case of  $n = 2$  has been worked out.  $\Leftarrow$  Arithmetic

# Power operations for Morava E-theory (height $n$ prime $p$ )

Theorem (Rezk '09, Barthel–Frankland '13)

If  $A = K(n)$ -local commutative  $E$ -algebra, then

$\pi_* A =$  graded amplified  $L$ -complete  $\Gamma$ -ring

- $\Gamma =$  twisted bialgebra over  $E_0$  (Dyer–Lashof algebra)
- $\exists Q_0 \in \Gamma$  with  $Q_0(x) \equiv x^p \pmod{p}$  (Frobenius congruence)

Goal make this structure explicit just as for Dyer–Lashof/Steenrod operations in ordinary homology.

The case of  $n = 2$  has been worked out.  $\Leftarrow$  Arithmetic

# Power operations for Morava E-theory (height $n$ prime $p$ )

Theorem (Rezk '09, Barthel–Frankland '13)

If  $A = K(n)$ -local commutative  $E$ -algebra, then

$\pi_* A =$  graded **amplified**  $L$ -complete  $\Gamma$ -ring

- $\Gamma =$  twisted bialgebra over  $E_0$  (Dyer–Lashof algebra)
- $\exists Q_0 \in \Gamma$  with  $Q_0(x) \equiv x^p \bmod p$  (Frobenius congruence)

Goal make this structure explicit just as for Dyer–Lashof/Steenrod operations in ordinary homology.

The case of  $n = 2$  has been worked out.  $\Leftarrow$  Arithmetic

# Power operations for Morava E-theory (height $n$ prime $p$ )

Theorem (Rezk '09, Barthel–Frankland '13)

If  $A = K(n)$ -local commutative  $E$ -algebra, then

$\pi_* A =$  graded amplified  **$L$ -complete**  $\Gamma$ -ring

- $\Gamma =$  twisted bialgebra over  $E_0$  (Dyer–Lashof algebra)
- $\exists Q_0 \in \Gamma$  with  $Q_0(x) \equiv x^p \pmod{p}$  (Frobenius congruence)

Goal make this structure explicit just as for Dyer–Lashof/Steenrod operations in ordinary homology.

The case of  $n = 2$  has been worked out.  $\Leftarrow$  Arithmetic

# Power operations for Morava E-theory (height $n$ prime $p$ )

Theorem (Rezk '09, Barthel–Frankland '13)

If  $A = K(n)$ -local commutative  $E$ -algebra, then

$\pi_* A =$  graded amplified  $L$ -complete  $\Gamma$ -ring

- $\Gamma =$  twisted bialgebra over  $E_0$  (Dyer–Lashof algebra)
- $\exists Q_0 \in \Gamma$  with  $Q_0(x) \equiv x^p \pmod{p}$  (Frobenius congruence)

Goal make this structure explicit just as for Dyer–Lashof/Steenrod operations in ordinary homology.

The case of  $n = 2$  has been worked out.  $\Leftarrow$  Arithmetic

# Power operations for Morava E-theory (height $n$ prime $p$ )

Theorem (Rezk '09, Barthel–Frankland '13)

If  $A = K(n)$ -local commutative  $E$ -algebra, then

$\pi_* A =$  graded amplified  $L$ -complete  $\Gamma$ -ring

- $\Gamma =$  twisted bialgebra over  $E_0$  (Dyer–Lashof algebra)
- $\exists Q_0 \in \Gamma$  with  $Q_0(x) \equiv x^p \pmod{p}$  (Frobenius congruence)

Goal make this structure explicit just as for Dyer–Lashof/Steenrod operations in ordinary homology.

The case of  $n = 2$  has been worked out.  $\Leftarrow$  Arithmetic

# Power operations for Morava E-theory (height $n$ prime $p$ )

Theorem (Rezk '09, Barthel–Frankland '13)

If  $A = K(n)$ -local commutative  $E$ -algebra, then

$\pi_* A =$  graded amplified  $L$ -complete  $\Gamma$ -ring

- $\Gamma =$  twisted bialgebra over  $E_0$  (Dyer–Lashof algebra)
- $\exists Q_0 \in \Gamma$  with  $Q_0(x) \equiv x^p \pmod{p}$  (Frobenius congruence)

Goal make this structure explicit just as for Dyer–Lashof/Steenrod operations in ordinary homology.

The case of  $n = 2$  has been worked out.  $\Leftarrow$  **Arithmetic**

# Power operations for Morava E-theory (height $n = 2$ )

## Theorem (Z. '19)

Given any Morava E-theory  $E$  of height 2 at a prime  $p$ , there is an explicit presentation for its algebra of power operations, in terms of generators  $Q_i: E^0(-) \rightarrow E^0(-)$ ,  $0 \leq i \leq p$ , and quadratic relations

$$Q_i Q_0 = - \sum_{k=1}^{p-i} w_0^k Q_{i+k} Q_k - \sum_{k=1}^p \sum_{m=0}^{k-1} w_0^m d_{i,k-m} Q_m Q_k$$

for  $1 \leq i \leq p$ , where the coefficients  $w_0$  and  $d_{i,k-m}$  arise from certain modular equations for elliptic curves.

Remark The first example, for  $p = 2$ , was calculated by Rezk '08. These have been applied to computations in unstable  $v_2$ -periodic homotopy theory (Z. '18 and ongoing joint work with G. Wang).



# Power operations for Morava E-theory (height $n = 2$ )

## Theorem (Z. '19)

Given any Morava E-theory  $E$  of height 2 at a prime  $p$ , there is an explicit presentation for its algebra of power operations, in terms of generators  $Q_i: E^0(-) \rightarrow E^0(-)$ ,  $0 \leq i \leq p$ , and quadratic relations

$$Q_i Q_0 = - \sum_{k=1}^{p-i} w_0^k Q_{i+k} Q_k - \sum_{k=1}^p \sum_{m=0}^{k-1} w_0^m d_{i,k-m} Q_m Q_k$$

for  $1 \leq i \leq p$ , where the coefficients  $w_0$  and  $d_{i,k-m}$  arise from certain modular equations for elliptic curves.

Remark The first example, for  $p = 2$ , was calculated by Rezk '08. These have been applied to computations in unstable  $v_2$ -periodic homotopy theory (Z. '18 and ongoing joint work with G. Wang).

# Power operations for Morava E-theory (height $n = 2$ )

## Theorem (Z. '19)

Given any Morava E-theory  $E$  of height 2 at a prime  $p$ , there is an explicit presentation for its algebra of power operations, in terms of generators  $Q_i: E^0(-) \rightarrow E^0(-)$ ,  $0 \leq i \leq p$ , and quadratic relations

$$Q_i Q_0 = - \sum_{k=1}^{p-i} w_0^k Q_{i+k} Q_k - \sum_{k=1}^p \sum_{m=0}^{k-1} w_0^m d_{i,k-m} Q_m Q_k$$

for  $1 \leq i \leq p$ , where the coefficients  $w_0$  and  $d_{i,k-m}$  arise from certain modular equations for elliptic curves.

Remark The first example, for  $p = 2$ , was calculated by Rezk '08. These have been applied to computations in unstable  $v_2$ -periodic homotopy theory (Z. '18 and ongoing joint work with G. Wang).

# Power operations for Morava E-theory (height $n = 2$ )

## Theorem (Z. '19)

Given any Morava E-theory  $E$  of height 2 at a prime  $p$ , there is an explicit presentation for its algebra of power operations, in terms of generators  $Q_i: E^0(-) \rightarrow E^0(-)$ ,  $0 \leq i \leq p$ , and quadratic relations

$$Q_i Q_0 = - \sum_{k=1}^{p-i} w_0^k Q_{i+k} Q_k - \sum_{k=1}^p \sum_{m=0}^{k-1} w_0^m d_{i,k-m} Q_m Q_k$$

for  $1 \leq i \leq p$ , where the coefficients  $w_0$  and  $d_{i,k-m}$  arise from certain **modular equations for elliptic curves**.

Remark The first example, for  $p = 2$ , was calculated by Rezk '08. These have been applied to computations in unstable  $v_2$ -periodic homotopy theory (Z. '18 and ongoing joint work with G. Wang).

# Power operations for Morava E-theory (height $n = 2$ )

## Theorem (Z. '19)

Given any Morava E-theory  $E$  of height 2 at a prime  $p$ , there is an explicit presentation for its algebra of power operations, in terms of generators  $Q_i: E^0(-) \rightarrow E^0(-)$ ,  $0 \leq i \leq p$ , and quadratic relations

$$Q_i Q_0 = - \sum_{k=1}^{p-i} w_0^k Q_{i+k} Q_k - \sum_{k=1}^p \sum_{m=0}^{k-1} w_0^m d_{i,k-m} Q_m Q_k$$

for  $1 \leq i \leq p$ , where the coefficients  $w_0$  and  $d_{i,k-m}$  arise from certain modular equations for elliptic curves.

Remark The first example, for  $p = 2$ , was calculated by Rezk '08. These have been applied to computations in unstable  $v_2$ -periodic homotopy theory (Z. '18 and ongoing joint work with G. Wang).

# Power operations for Morava E-theory (height $n = 2$ )

## Theorem (Z. '19)

Given any Morava E-theory  $E$  of height 2 at a prime  $p$ , there is an explicit presentation for its algebra of power operations, in terms of generators  $Q_i: E^0(-) \rightarrow E^0(-)$ ,  $0 \leq i \leq p$ , and quadratic relations

$$Q_i Q_0 = - \sum_{k=1}^{p-i} w_0^k Q_{i+k} Q_k - \sum_{k=1}^p \sum_{m=0}^{k-1} w_0^m d_{i,k-m} Q_m Q_k$$

for  $1 \leq i \leq p$ , where the coefficients  $w_0$  and  $d_{i,k-m}$  arise from certain modular equations for elliptic curves.

Remark The first example, for  $p = 2$ , was calculated by Rezk '08. These have been applied to computations in unstable  $v_2$ -periodic homotopy theory (Z. '18 and ongoing joint work with G. Wang).

# Power operations for Morava E-theory (height $n = 2$ )

## Theorem (Z. '19)

Given any Morava E-theory  $E$  of height 2 at a prime  $p$ , there is an explicit presentation for its algebra of power operations, in terms of generators  $Q_i: E^0(-) \rightarrow E^0(-)$ ,  $0 \leq i \leq p$ , and quadratic relations

$$Q_i Q_0 = - \sum_{k=1}^{p-i} w_0^k Q_{i+k} Q_k - \sum_{k=1}^p \sum_{m=0}^{k-1} w_0^m d_{i,k-m} Q_m Q_k$$

for  $1 \leq i \leq p$ , where the coefficients  $w_0$  and  $d_{i,k-m}$  arise from certain modular equations for elliptic curves.

Remark The first example, for  $p = 2$ , was calculated by Rezk '08. These have been applied to computations in unstable  $v_2$ -periodic homotopy theory (Z. '18 and ongoing joint work with G. Wang).

# Moduli of formal groups and algebras of power operations

Recall E-theory at height  $n$  and prime  $p$  has an underlying model

$$\begin{array}{ccc}
 F_k \xleftarrow{\text{univ defo}} \Gamma_{\mathbb{W}(k)}[[u_1, \dots, u_{n-1}]] & \longleftrightarrow & E \\
 \text{Frobenius isogenies} & & \text{power operations}
 \end{array}$$

An equivalence of cats (Ando–Hopkins–Strickland '04, Rezk '09)

$$\left\{ \begin{array}{l} \text{qcoh sheaves of grd comm algs} \\ \text{over the moduli problem of} \\ \text{defos of } F/k \text{ and Frob isogs} \end{array} \right\} \cong \left\{ \begin{array}{l} \text{grd comm algs over} \\ \text{the Dyer–Lashof algebra} \\ \text{for } E \end{array} \right\}$$

Goal Compute one side explicitly to get the other side.

# Moduli of formal groups and algebras of power operations

Recall E-theory at height  $n$  and prime  $p$  has an underlying model

$$\begin{array}{ccc}
 F_k & \xleftarrow{\text{univ defo}} & \Gamma_{\mathbb{W}(k)}[[u_1, \dots, u_{n-1}]] & \longleftrightarrow & E \\
 & & \text{Frobenius isogenies} & & \text{power operations}
 \end{array}$$

An equivalence of cats (Ando–Hopkins–Strickland '04, Rezk '09)

$$\left\{ \begin{array}{l} \text{qcoh sheaves of grd comm algs} \\ \text{over the moduli problem of} \\ \text{defos of } F/k \text{ and Frob isogs} \end{array} \right\} \cong \left\{ \begin{array}{l} \text{grd comm algs over} \\ \text{the Dyer–Lashof algebra} \\ \text{for } E \end{array} \right\}$$

Goal Compute one side explicitly to get the other side.



# Moduli of formal groups and algebras of power operations

Recall E-theory at height  $n$  and prime  $p$  has an underlying model

$$\begin{array}{ccc}
 F_k & \xleftarrow{\text{univ defo}} & \Gamma_{\mathbb{W}(k)}[[u_1, \dots, u_{n-1}]] & \longleftrightarrow & E \\
 & & \text{Frobenius isogenies} & & \text{power operations}
 \end{array}$$

An equivalence of cats (Ando–Hopkins–Strickland '04, Rezk '09)

$$\left\{ \begin{array}{l} \text{qcoh sheaves of grd comm algs} \\ \text{over the moduli problem of} \\ \text{defos of } F/k \text{ and Frob isogs} \end{array} \right\} \cong \left\{ \begin{array}{l} \text{grd comm algs over} \\ \text{the Dyer–Lashof algebra} \\ \text{for } E \end{array} \right\}$$

Goal Compute one side explicitly to get the other side.

# Moduli of formal groups and algebras of power operations

Recall E-theory at height  $n$  and prime  $p$  has an underlying model

$$\begin{array}{ccc}
 F_k \xleftarrow{\text{univ defo}} \Gamma_{\mathbb{W}(k)}[[u_1, \dots, u_{n-1}]] & \longleftrightarrow & E \\
 \text{Frobenius isogenies} & & \text{power operations}
 \end{array}$$

An equivalence of cats (Ando–Hopkins–Strickland '04, Rezk '09)

$$\left\{ \begin{array}{l} \text{qcoh sheaves of grd comm algs} \\ \text{over the moduli problem of} \\ \text{defos of } F/k \text{ and Frob isogs} \end{array} \right\} \cong \left\{ \begin{array}{l} \text{grd comm algs over} \\ \text{the Dyer–Lashof algebra} \\ \text{for } E \end{array} \right\}$$

Goal Compute one side explicitly to get the other side.

# Moduli of formal groups and algebras of power operations

Recall E-theory at height  $n$  and prime  $p$  has an underlying model

$$\begin{array}{ccc} F_k \xleftarrow{\text{univ defo}} \Gamma_{\mathbb{W}(k)}[[u_1, \dots, u_{n-1}]] & \longleftrightarrow & E \\ \text{Frobenius isogenies} & & \text{power operations} \end{array}$$

An equivalence of cats (Ando–Hopkins–Strickland '04, Rezk '09)

$$\left\{ \begin{array}{l} \text{qcoh sheaves of grd comm algs} \\ \text{over the moduli problem of} \\ \text{defos of } F/k \text{ and Frob isogs} \end{array} \right\} \cong \left\{ \begin{array}{l} \text{grd comm algs over} \\ \text{the Dyer–Lashof algebra} \\ \text{for } E \end{array} \right\}$$

Goal Compute one side explicitly to get the other side.

# Moduli of formal groups and algebras of power operations

Recall E-theory at height  $n$  and prime  $p$  has an underlying model

$$\begin{array}{ccc} F_k & \xleftarrow{\text{univ defo}} & \Gamma_{\mathbb{W}(k)}[[u_1, \dots, u_{n-1}]] \\ & \circlearrowleft & \\ & \text{Frobenius isogenies} & \end{array} \quad \begin{array}{ccc} & \longleftrightarrow & E \\ & & \circlearrowleft \\ & \text{power operations} & \end{array}$$

An equivalence of cats (Ando–Hopkins–Strickland '04, Rezk '09)

$$\left\{ \begin{array}{l} \text{qcoh sheaves of grd comm algs} \\ \text{over the moduli problem of} \\ \text{defos of } F/k \text{ and Frob isogs} \end{array} \right\} \cong \left\{ \begin{array}{l} \text{grd comm algs over} \\ \text{the Dyer–Lashof algebra} \\ \text{for } E \end{array} \right\}$$

Goal Compute one side explicitly to get the other side.

# Moduli of formal groups and algebras of power operations

Recall E-theory at height  $n$  and prime  $p$  has an underlying model

$$\begin{array}{ccc}
 F_k & \xleftarrow{\text{univ defo}} & \Gamma_{\mathbb{W}(k)}[[u_1, \dots, u_{n-1}]] & \longleftrightarrow & E \\
 & & \text{Frobenius isogenies} & & \text{power operations}
 \end{array}$$

An equivalence of cats (Ando–Hopkins–Strickland '04, Rezk '09)

$$\left\{ \begin{array}{l} \text{qcoh sheaves of grd comm algs} \\ \text{over the moduli problem of} \\ \text{defos of } F/k \text{ and Frob isogs} \end{array} \right\} \cong \left\{ \begin{array}{l} \text{grd comm algs over} \\ \text{the Dyer–Lashof algebra} \\ \text{for } E \end{array} \right\}$$

Goal Compute one side explicitly to get the other side.

# Moduli of formal groups and algebras of power operations

Recall E-theory at height  $n$  and prime  $p$  has an underlying model

$$\begin{array}{ccc} F_k & \xleftarrow{\text{univ defo}} & \Gamma_{\mathbb{W}(k)}[[u_1, \dots, u_{n-1}]] \\ & \circlearrowleft & \\ & \text{Frobenius isogenies} & \end{array} \quad \begin{array}{ccc} & \longleftrightarrow & E \\ & & \circlearrowleft \\ & \text{power operations} & \end{array}$$

An equivalence of cats (Ando–Hopkins–Strickland '04, Rezk '09)

$$\left\{ \begin{array}{l} \text{qcoh sheaves of grd comm algs} \\ \text{over the moduli problem of} \\ \text{defos of } F/k \text{ and Frob isogs} \end{array} \right\} \cong \left\{ \begin{array}{l} \text{grd comm algs over} \\ \text{the Dyer–Lashof algebra} \\ \text{for } E \end{array} \right\}$$

Goal Compute one side explicitly to get the other side.

# Moduli of formal groups and algebras of power operations

Recall E-theory at height  $n$  and prime  $p$  has an underlying model

$$\begin{array}{ccc} F_k & \xleftarrow{\text{univ defo}} & \Gamma_{\mathbb{W}(k)}[[u_1, \dots, u_{n-1}]] \\ & \circlearrowleft & \\ & \text{Frobenius isogenies} & \end{array} \quad \begin{array}{ccc} & \longleftrightarrow & E \\ & & \circlearrowleft \\ & & \text{power operations} \end{array}$$

An equivalence of cats (Ando–Hopkins–Strickland '04, Rezk '09)

$$\left\{ \begin{array}{l} \text{qcoh sheaves of grd comm algs} \\ \text{over the moduli problem of} \\ \text{defos of } F/k \text{ and Frob isogs} \end{array} \right\} \cong \left\{ \begin{array}{l} \text{grd comm algs over} \\ \text{the Dyer–Lashof algebra} \\ \text{for } E \end{array} \right\}$$

Goal Compute one side explicitly to get the other side.

# Moduli of formal groups and algebras of power operations

Recall E-theory at height  $n$  and prime  $p$  has an underlying model

$$\begin{array}{ccc} F_k & \xleftarrow{\text{univ defo}} & \Gamma_{\mathbb{W}(k)}[[u_1, \dots, u_{n-1}]] \\ & \circlearrowleft & \\ & \text{Frobenius isogenies} & \end{array} \quad \longleftrightarrow \quad \begin{array}{ccc} E & & \\ & \circlearrowleft & \\ & \text{power operations} & \end{array}$$

An equivalence of cats (Ando–Hopkins–Strickland '04, Rezk '09)



$$\left\{ \begin{array}{l} \text{qcoh sheaves of grd comm algs} \\ \text{over the moduli problem of} \\ \text{defos of } F/k \text{ and Frob isogs} \end{array} \right\} \cong \left\{ \begin{array}{l} \text{grd comm algs over} \\ \text{the Dyer–Lashof algebra} \\ \text{for } E \end{array} \right\}$$

Goal Compute one side explicitly to get the other side.



# Moduli of elliptic curves and D.-L. algebras at height 2

Moduli of formal groups and moduli of ell. curves (Serre–Tate '64)  
 $p$ -adically, defo thy of an ec  $\cong$  defo thy of its  $p$ -divisible gp  
 $[\Gamma_0(p)]$  as an *open arithmetic surface* (Katz–Mazur '85)  
parameters for its local ring at a supersingular point, chosen from  
specific modular forms





- Compactify the moduli
- Compute with explicit  $q$ -expansions
- Transport from cusps to s.sing. pt

## Theorem (Z. '19)

A choice of such parameters,  $h$  and  $\alpha$ , satisfies the equation  
$$(\alpha - 1)^p(\alpha - p) - ((-1)^p + (-1)^{p-1}(-p)p + h)\alpha = 0$$

# Moduli of elliptic curves and D.-L. algebras at height 2

Moduli of formal groups and moduli of ell. curves (Serre–Tate '64)  
 $p$ -adically, defo thy of an ec  $\cong$  defo thy of its  $p$ -divisible gp  
 $[\Gamma_0(p)]$  as an *open arithmetic surface* (Katz–Mazur '85)  
parameters for its local ring at a supersingular point, chosen from  
specific modular forms





- Compactify the moduli
- Compute with explicit  $q$ -expansions
- Transport from cusps to s.sing. pt

## Theorem (Z. '19)

A choice of such parameters,  $h$  and  $\alpha$ , satisfies the equation  
$$(\alpha - 1)^p(\alpha - p) - ((-1)^p + (-1)^{p-1}(-p)p + h)\alpha = 0$$

# Moduli of elliptic curves and D.-L. algebras at height 2

Moduli of formal groups and moduli of ell. curves (Serre–Tate '64)  
 $p$ -adically, defo thy of a **s.sing.** ec  $\cong$  defo thy of its  $p$ -divisible gp  
 $[\Gamma_0(p)]$  as an *open arithmetic surface* (Katz–Mazur '85)  
parameters for its local ring at a supersingular point, chosen from  
specific modular forms





- Compactify the moduli
- Compute with explicit  $q$ -expansions
- Transport from cusps to s.sing. pt

## Theorem (Z. '19)

A choice of such parameters,  $h$  and  $\alpha$ , satisfies the equation  
$$(\alpha - 1)^p(\alpha - p) - ((-1)^p + (-1)^{p-1}(-p)p + h)\alpha = 0$$

# Moduli of elliptic curves and D.-L. algebras at height 2

Moduli of formal groups and moduli of ell. curves (Serre–Tate '64)  
 $p$ -adically, defo thy of a **s.sing.** ec  $\cong$  defo thy of its **formal** gp  
 $[\Gamma_0(p)]$  as an *open arithmetic surface* (Katz–Mazur '85)  
parameters for its local ring at a supersingular point, chosen from  
specific modular forms





- Compactify the moduli
- Compute with explicit  $q$ -expansions
- Transport from cusps to s.sing. pt

## Theorem (Z. '19)

A choice of such parameters,  $h$  and  $\alpha$ , satisfies the equation  
$$(\alpha - 1)^p(\alpha - p) - ((-1)^p + (-1)^{p-1}(-p)p + h)\alpha = 0$$

# Moduli of elliptic curves and D.-L. algebras at height 2

Moduli of formal groups and moduli of ell. curves (Serre–Tate '64)  
 $p$ -adically, defo thy of a s.sing. ec  $\cong$  defo thy of its formal gp  
 $[\Gamma_0(p)]$  as an *open arithmetic surface* (Katz–Mazur '85)  
parameters for its local ring at a supersingular point, chosen from  
specific modular forms





- Compactify the moduli
- Compute with explicit  $q$ -expansions
- Transport from cusps to s.sing. pt

## Theorem (Z. '19)

A choice of such parameters,  $h$  and  $\alpha$ , satisfies the equation  
$$(\alpha - 1)^p(\alpha - p) - ((-1)^p + (-1)^{p-1}(-p)p + h)\alpha = 0$$

# Moduli of elliptic curves and D.-L. algebras at height 2

Moduli of formal groups and moduli of ell. curves (Serre–Tate '64)  
 $p$ -adically, defo thy of a s.sing. ec  $\cong$  defo thy of its formal gp  
 $[\Gamma_0(p)]$  as an *open arithmetic surface* (Katz–Mazur '85)  
parameters for its local ring at a supersingular point, chosen from  
specific modular forms





- Compactify the moduli
- Compute with explicit  $q$ -expansions
- Transport from cusps to s.sing. pt

## Theorem (Z. '19)

A choice of such parameters,  $h$  and  $\alpha$ , satisfies the equation  
$$(\alpha - 1)^p(\alpha - p) - ((-1)^p + (-1)^{p-1}(-p)p + h)\alpha = 0$$

# Moduli of elliptic curves and D.-L. algebras at height 2

Moduli of formal groups and moduli of ell. curves (Serre–Tate '64)  
 $p$ -adically, defo thy of a s.sing. ec  $\cong$  defo thy of its formal gp  
 $[\Gamma_0(p)]$  as an *open arithmetic surface* (Katz–Mazur '85)  
parameters for its local ring at a supersingular point, chosen from  
specific modular forms





- **Compactify** the moduli
- Compute with explicit  $q$ -expansions
- Transport from cusps to s.sing. pt

## Theorem (Z. '19)

A choice of such parameters,  $h$  and  $\alpha$ , satisfies the equation  
$$(\alpha - 1)^p(\alpha - p) - ((-1)^p + (-1)^{p-1}(-p)p + h)\alpha = 0$$

# Moduli of elliptic curves and D.-L. algebras at height 2

Moduli of formal groups and moduli of ell. curves (Serre–Tate '64)  
 $p$ -adically, defo thy of a s.sing. ec  $\cong$  defo thy of its formal gp  
 $[\Gamma_0(p)]$  as an *open arithmetic surface* (Katz–Mazur '85)  
parameters for its local ring at a supersingular point, chosen from  
**specific modular forms**



- Compactify the moduli
- Compute with **explicit  $q$ -expansions**
- Transport from cusps to s.sing. pt



## Theorem (Z. '19)

A choice of such parameters,  $h$  and  $\alpha$ , satisfies the equation  
$$(\alpha - 1)^p(\alpha - p) - ((-1)^p + (-1)^{p-1}(-p)p + h)\alpha = 0$$



# Moduli of elliptic curves and D.-L. algebras at height 2

Moduli of formal groups and moduli of ell. curves (Serre–Tate '64)  
 $p$ -adically, defo thy of a s.sing. ec  $\cong$  defo thy of its formal gp  
 $[\Gamma_0(p)]$  as an *open arithmetic surface* (Katz–Mazur '85)  
parameters for its local ring at a supersingular point, chosen from  
specific modular forms





- Compactify the moduli
- Compute with explicit  $q$ -expansions
- Transport from cusps to s.sing. pt

## Theorem (Z. '19)

A choice of such parameters,  $h$  and  $\alpha$ , satisfies the equation  
$$(\alpha - 1)^p(\alpha - p) - ((-1)^p + (-1)^{p-1}(-p)p + h)\alpha = 0$$

# Moduli of elliptic curves and D.-L. algebras at height 2

Moduli of formal groups and moduli of ell. curves (Serre–Tate '64)  
 $p$ -adically, defo thy of a s.sing. ec  $\cong$  defo thy of its formal gp  
 $[\Gamma_0(p)]$  as an *open arithmetic surface* (Katz–Mazur '85)  
parameters for its local ring at a supersingular point, chosen from  
specific modular forms





- Compactify the moduli
- Compute with explicit  $q$ -expansions
- Transport from cusps to s.sing. pt

## Theorem (Z. '19)

A choice of such parameters,  $h$  and  $\alpha$ , satisfies the equation  
$$(\alpha - 1)^p(\alpha - p) - ((-1)^p + (-1)^{p-1}(-p)p + h)\alpha = 0$$

# Moduli of elliptic curves and D.-L. algebras at height 2

Moduli of formal groups and moduli of ell. curves (Serre–Tate '64)  
 $p$ -adically, defo thy of a s.sing. ec  $\cong$  defo thy of its formal gp  
 $[\Gamma_0(p)]$  as an *open arithmetic surface* (Katz–Mazur '85)  
parameters for its local ring at a supersingular point, chosen from  
specific modular forms





- Compactify the moduli
- Compute with explicit  $q$ -expansions
- Transport from **cusps** to s.sing. pt

## Theorem (Z. '19)

A choice of such parameters,  $h$  and  $\alpha$ , satisfies the equation  
 $(\alpha - 1)^p(\alpha - p) - ((-1)^p + (-1)^{p-1}(-p)p + h)\alpha = 0$

# Moduli of elliptic curves and D.-L. algebras at height 2

Moduli of formal groups and moduli of ell. curves (Serre–Tate '64)  
 $p$ -adically, defo thy of a s.sing. ec  $\cong$  defo thy of its formal gp  
 $[\Gamma_0(p)]$  as an *open arithmetic surface* (Katz–Mazur '85)  
parameters for its local ring at a supersingular point, chosen from  
specific modular forms





- Compactify the moduli
- Compute with explicit  $q$ -expansions
- Transport from cusps to s.sing. pt

## Theorem (Z. '19)

A choice of such parameters,  $h$  and  $\alpha$ , satisfies the equation  
$$(\alpha - 1)^p(\alpha - p) - ((-1)^p + (-1)^{p-1}(-p)p + h)\alpha = 0$$

# Moduli of elliptic curves and D.-L. algebras at height 2

Moduli of formal groups and moduli of ell. curves (Serre–Tate '64)  
 $p$ -adically, defo thy of a s.sing. ec  $\cong$  defo thy of its formal gp  
 $[\Gamma_0(p)]$  as an *open arithmetic surface* (Katz–Mazur '85)  
parameters for its local ring at a supersingular point, chosen from  
specific modular forms





- Compactify the moduli
- Compute with explicit  $q$ -expansions
- Transport from cusps to s.sing. pt

## Theorem (Z. '19)

A choice of such parameters,  $h$  and  $\alpha$ , satisfies the equation  
 $(\alpha - 1)^p(\alpha - p) - ((-1)^p + (-1)^{p-1}(-p)p + \textcolor{red}{h})\alpha = 0$

# Moduli of elliptic curves and D.-L. algebras at height 2

Moduli of formal groups and moduli of ell. curves (Serre–Tate '64)  
 $p$ -adically, defo thy of a s.sing. ec  $\cong$  defo thy of its formal gp  
 $[\Gamma_0(p)]$  as an *open arithmetic surface* (Katz–Mazur '85)  
parameters for its local ring at a supersingular point, chosen from  
specific modular forms





- Compactify the moduli
- Compute with explicit  $q$ -expansions
- Transport from cusps to s.sing. pt

## Theorem (Z. '19)

A choice of such parameters,  $h$  and  $\alpha$ , satisfies the equation  
 $(\alpha - 1)^p(\alpha - p) - ((-1)^p + (-1)^{p-1}(-p)p + h)\alpha \equiv \alpha(\alpha^p - h) \pmod{p}$

# Moduli of elliptic curves and D.-L. algebras at height 2

Moduli of formal groups and moduli of ell. curves (Serre–Tate '64)  
 $p$ -adically, defo thy of a s.sing. ec  $\cong$  defo thy of its formal gp  
 $[\Gamma_0(p)]$  as an *open arithmetic surface* (Katz–Mazur '85)  
parameters for its local ring at a supersingular point, chosen from  
specific modular forms





- Compactify the moduli
- Compute with explicit  $q$ -expansions
- Transport from cusps to s.sing. pt

## Theorem (Z. '19)

A choice of such parameters,  $h$  and  $\alpha$ , satisfies the equation  
 $(\alpha - 1)^p(\alpha - p) - ((-1)^p + (-1)^{p-1}(-p)p + h)\alpha \equiv \alpha(\tilde{j}^p - j) \pmod{p}$

# Moduli of elliptic curves and D.-L. algebras at height 2

Moduli of formal groups and moduli of ell. curves (Serre–Tate '64)  
 $p$ -adically, defo thy of a s.sing. ec  $\cong$  defo thy of its formal gp  
 $[\Gamma_0(p)]$  as an *open arithmetic surface* (Katz–Mazur '85)  
parameters for its local ring at a supersingular point, chosen from  
specific modular forms



- Compactify the moduli
- Compute with explicit  $q$ -expansions
- Transport from cusps to s.sing. pt

## Theorem (Z. '19)

A choice of such parameters,  $h$  and  $\alpha$ , satisfies the equation  
$$(\alpha - 1)^p(\alpha - p) - ((-1)^p + (-1)^{p-1}(-p)p + h)\alpha \equiv \alpha(\alpha^p - h) \pmod{p}$$




# Moduli of elliptic curves and D.-L. algebras at height 2

Question At height  $n > 2$ , can we get an explicit presentation for the Dyer–Lashof algebra of Morava E-theory?

Investigating J. Weinstein’s approach to integral models for modular curves via the *infinite* Lubin–Tate tower.

A picture from Jared Weinstein, Semistable models for modular curves of arbitrary level




# Moduli of elliptic curves and D.-L. algebras at height 2

Question At height  $n > 2$ , can we get an explicit presentation for the Dyer–Lashof algebra of Morava E-theory?

Investigating J. Weinstein’s approach to integral models for modular curves via the *infinite* Lubin–Tate tower.

A picture from Jared Weinstein, Semistable models for modular curves of arbitrary level




# Moduli of elliptic curves and D.-L. algebras at height 2

Question At height  $n > 2$ , can we get an explicit presentation for the Dyer–Lashof algebra of Morava E-theory?

Investigating J. Weinstein’s approach to integral models for modular curves via the *infinite* Lubin–Tate tower.

A picture from Jared Weinstein, Semistable models for modular curves of arbitrary level



- Established in 2012, a public research-oriented university funded by the municipal government of Shenzhen, China's innovation center.
- Over 800 faculty members, 4205 undergrad students, 2214 postgrad/doctoral students, international students from 15 countries, student:teacher = 10:1.
- Set on five hundred acres of wooded landscape in the picturesque South Mountain area.
- Department of Mathematics founded in June 2015.  
28 research-and-teaching-line faculty members, 6 teaching-line faculty members, 101 grad students, 205 undergrad majors.
- International Center for Mathematics founded in February 2019.

*Thank you.*